

Flavor physics at Future Circular Lepton Collider

1000,000,000,000+


As a Tera - Z +

The text "As a Tera - Z +" is centered. To the right of the "Z" is a large blue plus sign. There are two blue circles with light blue conical shadows: one is positioned above the plus sign and the other is below the "Z".

Lingfeng Li, Brown U.

Sep. 12, 2022

ICFA Advanced Beam Dynamics Workshop on High Luminosity Circular e+e- Colliders



Prologue

“Don’t just leave flavor physics to flavor physicists.”


[Someone Awesome, 2019?]

“Non-flavor physicists must be amused first.”

[me, 2022]

Tera-Z as a Z and Flavor Factory


b -hadrons	Belle II ($50+5 \text{ ab}^{-1}$)	LHCb (300 fb^{-1})	Tera-Z
B^0, \bar{B}^0	5.4×10^{10} (50 ab^{-1} on $\Upsilon(4S)$)	3×10^{13}	1.2×10^{11}
B^\pm	5.7×10^{10} (50 ab^{-1} on $\Upsilon(4S)$)	3×10^{13}	1.2×10^{11}
B_s^0, \bar{B}_s^0	6.0×10^8 (5 ab^{-1} on $\Upsilon(5S)$)	1×10^{13}	3.1×10^{10}
B_c^\pm	-	1×10^{11}	1.8×10^8
$\Lambda_b^0, \bar{\Lambda}_b^0$	-	2×10^{13}	2.5×10^{10}
$c(\bar{c})$	2.6×10^{11}	$\gtrsim 10^{14}$	2.4×10^{11}
τ^\pm	9×10^{10}	-	7.4×10^{10}



- Higher luminosity as the accelerator design keeps upgrading
- ≥ 2 interaction points and various detectors

Flavor physics also need energy larger than 91 GeV (e.g., $|V_{cb}|$ from W decays)

~100/ π km





Still a lot to understand
even we can write down \mathcal{L}_{SM}



Great ways to probe new physics
Great prize awaits?



Who Ordered These?

Flavor mixing and CP violation patterns


Large flavor hierarchy




Flavor and New Physics

Heavy flavors (b, c, and τ) are long-lived particles, width $< 10^{-11}$ GeV \ll mass:

$$\Gamma_{\text{SM}} \sim \frac{G_F^2 m_f^5}{192\pi^3} \times \text{const} \propto \frac{m_f^5}{m_W^4}.$$




$$\Gamma_{\text{BSM}} \propto \frac{m_f^5}{\Lambda_{\text{NP}}^2 m_W^2} \text{ (w/ interference), or } \frac{m_f^5}{\Lambda_{\text{NP}}^4} \text{ (w/o interference)}$$




$$\left(\frac{m_W^2}{\Lambda_{\text{NP}}^2} \text{ or } \frac{m_W^4}{\Lambda_{\text{NP}}^4} \gg \frac{m_f^4}{\Lambda_{\text{NP}}^4} \right)$$

Large rates with moderate suppression



Recognizing “Golden” Modes

- Neutrinos
- Neutrals
(photon/ π^0 / η ...)
- Rare modes
- τ decays
- BSM states



- Baryonic tracks
- Electron and Muon
- $b \rightarrow c \rightarrow \tau$ cascade
- Long-lived particles
- Boost: O(fs) time scales
- Heavy species: Bc, Λ_b , tetraquarks...
- Multiple soft tracks

Recent Progress

Disclaimer: *Priorities are given to numerical results with (fast or full) simulations in stead of theoretical works.*

Apologize for any missing contributions due to personal ignorance and prejudice.

Precision Flavor Measurements


[J. Charles, Z. Ligeti, S. Monteil, M. Papucci et al., 2006.04824](#)

	Central values	Current [18]	Uncertainties		+FCC-ee
			Phase I	Phase II	Phase III
$ V_{ud} $	0.97437	± 0.00021	id	id	id
$ V_{us} f_+^{K \rightarrow \pi}(0)$	0.2177	± 0.0004	id	id	id
$ V_{cd} $	0.2248	± 0.0043	± 0.003	id	id
$ V_{cs} $	0.9735	± 0.0094	id	id	id
Δm_d [ps ⁻¹]	0.5065	± 0.0019	id	id	id
Δm_s [ps ⁻¹]	17.757	± 0.021	id	id	id
$ V_{cb} _{\text{SL}} \times 10^3$	42.26	± 0.58	± 0.60	± 0.44	id
$ V_{cb} _{W \rightarrow cb} \times 10^3$	—	—	—	—	± 0.17
$ V_{ub} _{\text{SL}} \times 10^3$	3.56	± 0.22	± 0.042	± 0.032	id
$ V_{ub}/V_{cb} $ (from Λ_b)	0.0842	± 0.0050	± 0.0025	± 0.0008	id
$\mathcal{B}(B \rightarrow \tau \nu) \times 10^4$	0.83	± 0.24	± 0.04	± 0.02	± 0.009
$\mathcal{B}(B \rightarrow \mu \nu) \times 10^6$	0.37	—	± 0.03	± 0.02	id
$\sin 2\beta$	0.680	± 0.017	± 0.005	± 0.002	± 0.0008
α [°] (mod 180°)	91.9	± 4.4	± 0.6	id	id
γ [°] (mod 180°)	66.7	± 5.6	± 1	± 0.25	± 0.20
β_s [rad]	-0.035	± 0.021	± 0.014	± 0.004	± 0.002
$A_{\text{SL}}^d \times 10^4$	-6	± 19	± 5	± 2	± 0.25
$A_{\text{SL}}^s \times 10^5$	3	± 300	± 70	± 30	± 2.5

Lot's of unknown values (identical to previous phases) in the Tera-Z era
Expect improvements?

Also many values to be verified (may need more simulations)
Both stat. and syst.

Precision Flavor Measurements (II)




Probing high BSM scales with $\Delta F=2$ measurements (future Belle II + LHCb)

See Luca Selvestrini's talk & [J. Charles, Z. Ligeti, S. Monteil, M. Papucci et al., 2006.04824](https://arxiv.org/abs/2006.04824) for more details

CPV Angles


- Multiple new ways of measurement
- Current focus: B decays

See also: [J. Charles, S. Descotes-Genon, Zoltan Ligeti, S. Monteil, M. Papucci, K. Trabelsi, L. Silva, 2006.04824](#)
[R. Aleksan, L. Oliver, E. Perez, 2107.05311](#)
[X. Li, M Ruan, M. Zhao, 2205.10565](#)




⬆: Measurement of α using $B \rightarrow \pi^0 \pi^0 \rightarrow 4\gamma$ (!)
 Removing mirror solutions

[Y. Wang, S. Descotes-Genon, O. Deschamps, LL, S. Chen, Y. Zhu, M. Ruan, 2208.08237](#)




FCNC: Dileptonic Modes

- Rare decays, sensitive to BSM
- Partially motivated by R_K and R_{K^*} anomalies
- Flagship mode: $b \rightarrow s \tau \tau$, highly sensitive to LFUV in 3rd generation



[LL, T. Liu, 2012.00665](#)

Probing rare decay BR @ $O(10^{-7})$ level*
*: Absolute BR




⬆: Even better background mitigation by π^0 reconstruction from background D_s decays

See also:

[J. F. Kamenik, S. Monteil, A. Semkiv, L. V. Silva, 1705. 11106](#) **14**


FCNC: Dileptonic Modes (II)




→ : Extended to flavor violating modes, e.g., $B_d \rightarrow K^* \tau \mu$.

Lingfeng Li, Brown U.

← : Measurements for both $B_d \rightarrow \mu \mu$ & $B_s \rightarrow \mu \mu$. $B_d \rightarrow \pi \pi$ background under control due to advanced PID tech.



FCNC: Di-neutrino Modes & More




☐←: Reconstruct $b \rightarrow sv\nu$ semi-invisible decay, error on $q^2 < 3 \text{ GeV}^2$



- Can also constraint BSM physics, e.g., axion(-(like-particle)) from flavor decays [J. Camalich, M. Pospelov, P. Vuong, R. Ziegler, J. Zupan, 2002.04623](#)
- Great potential for radiative decays

Heavy Quark Weak Decays (FCCC)

[T. Zheng, J. Xu, L. Cao, D. Yu, W. Wang et al., 2007.08234](#)





- Anomalies indicating lepton flavor universality violation
- Potential for $|V_{cb}|$ & $|V_{ub}|$ extraction
- Current focus: (Semi)leptonic modes



⬆: $R_{J/\psi}$, $R_{D_s^{(*)}}$, R_{Λ_b} projections on the way

[T. Kwok, X. Jiang, LL, Tao Liu, In prep](#)




Probe O(10) TeV BSM in 9 dimensions by combining multiple semileptonic & rare modes

Tau and Lepton Sector

[L. Calibbi, T. Li, X. Marcano, M.A. Schmidt, 2207.10913](#)

- A most powerful tau machine
- Current focus: charged lepton flavor violation (cLFV)



↗: Complementarity on cLFV new physics between exotic Z, quarkonia, and lepton decays


See also: [L. Calibbi, X. Marcano, J. Roy, 2107.10273](#)

Tau and Lepton Sector (II)

Interesting studies include:


- Lifetime (better vertex resolution)
- Lepton universality via tau decays (good lepton PID)
- Hadronic decays and QCD coupling (ECAL resolution)
- Polarimetry (also for EW)

More discussions to be found in Alberto Lusiani's talk



[M. Dam, 1811.09408](#)

[M. Dam, 2107.12832](#)



Summary: A flavor-centric perspective

- Origin of matter?
understand lepton and baryon numbers
- Light dark matter?
- Lepton Flavor
Universality anomalies?

BSM



Higgs

EWPT
Top

QCD

Hardware

- Origin of flavor hierarchy?
- CP violation phases from Yukawa?

- Flavor physics beyond the Tera-Z phase?
- Common need in τ phys.

- How does asymptotic freedom work with flavor?
- New formalism beyond the conventional meson-baryon picture?

- Use a plethora of data to improve hadronization

Most demanding field:
We need better tracker, E(H)CAL, electronics... everything!